


The Role of Food in Maintaining Immune Health in Ageing

Fiona McEvoy Ph.D. & Christine Loscher Ph.D.

Part 2: How Does the Immune System Work? A Deep Dive into the Innate and Adaptive Immune System



What Does the Immune System do?

The immune system is a collection of cells, tissues and organs that has evolved to defend against dangerous pathogens and maintain homeostasis within an organism. The immune system maintains a constant surveillance against harmful bacteria and viruses, and has the ability to combat both foreign invaders and abnormal endogenous factors such as cancerous cells. There are two branches of the immune system broadly divided into the innate and adaptive systems ⁴. The innate immune system represents the rapid first line of defence against invading pathogens and is present in all multicellular organisms. Initial protection is provided by physical barriers to entry such as the skin and epithelial linings of the respiratory, gastrointestinal and mucosal systems. If a breach of these barriers occurs, cells of the innate immune system including monocytes, macrophages and neutrophils attempt to prevent invasion by means of inducing inflammation, phagocytosis of pathogens, release of lytic enzymes and activation of complement pathways ⁵.

During these processes, dendritic cells (also known as antigen-presenting cells) engulf a portion of the antigen and migrate to the lymph nodes where they interact with, and activate T lymphocytes to initiate an adaptive immune response. The adaptive immune response is antigen specific and can initially take between four and seven days to develop. However, due to immunological memory, subsequent exposure to the same pathogen results in much quicker immune responses. There are a number of different T cell populations whose activation depends on the type of pathogen encountered ⁶. CD8+ cytotoxic T cells are very important for immune defence against intracellular pathogens and for tumour surveillance. CD4+ T helper cells (of which there are many varieties) play a major role in instigating and shaping adaptive immune responses via the secretion of various cytokines ⁷. Furthermore, B lymphocytes (B cells) are responsible for producing antibodies that target extracellular pathogens.

The Innate Immune System

Neutrophils

Neutrophils are the first white blood cells recruited to sites of acute inflammation, in response to chemotactic cues such as IL-8, produced by stressed tissue cells and tissue-resident macrophages. Neutrophils constitute the primary immune defence against rapidly dividing bacteria, yeast and fungal infections, deploying microbicidal (i.e. destroys microbes) mechanisms including the generation of reactive oxygen and nitrogen species and the release of proteolytic enzymes and microbicidal peptides. The appropriate initiation and resolution of their inflammatory responses is crucial to the clearance of infections and the prevention of non-specific tissue damage leading to chronic inflammatory disease.

Neutrophils constitute the largest population of white blood cells in humans and mice, and to date there has been no age-related decline in numbers observed ¹⁴. However, neutrophils do show an age-related decline in most aspects of their microbicidal functions including chemotaxis, phagocytosis of microbes and generation of superoxide in response to stimulation by soluble host and bacterial factors ^{15,16}. The reductions in chemotactic and phagocytic ability observed in neutrophils from older donors would result in a delayed neutrophilic response to bacterial invasion, allowing rapidly dividing bacteria to establish a stronger core of infection. Inefficient chemotaxis also has the ability to increase tissue damage because neutrophils secrete proteases such as elastase to aid their migration through tissues; this is thought to extend inflammation and impair resolution of inflammation in older adults ¹⁴. Neutrophils are the shortest-lived blood cells and their life span is extended at sites of infection by survival signals provided by cytokines (e.g. GM-CSF) and bacterial products (e.g. LPS). The age-associated reduction in neutrophils' ability to respond to survival signals, specifically GM-CSF, will result in compromised removal of microbes and thus extend the time to resolve the inflammation ¹⁷.

Monocytes / Macrophage

Monocytes originate from myeloid stem cell progenitors and differentiate into macrophages with specialised functions depending on their location within the body ¹⁸. Macrophages are specialised cells involved in the detection and destruction of bacteria and other harmful pathogens via phagocytosis and production of reactive oxygen and nitrogen species. They are also capable of releasing a vast range of cytokines and chemokines that are essential for the initiation and propagation of the inflammatory process ¹⁹. Macrophages are able to detect products of bacteria and other microorganisms using a system of pattern recognition receptors known as Toll-like receptors (TLRs). The TLRs are a family of invariant pattern recognition receptors with specificity for highly conserved portions of pathogens; 11 human TLRs have been identified to date. TLR activation results in both proinflammatory cytokine response via NF- κ B-dependant pathway and the upregulation of type I IFN and IFN dependant genes ⁵.

There is conflicting data surrounding the effect of ageing on macrophage chemotaxis and phagocytosis with some studies showing decreases while others report the opposite ²⁰. There is now however a large body of data to support the theory that ageing alters cytokine secretion by macrophages in response to TLR stimulation. Age-associated decreases in LPS-induced (activating TLR4) IL-1 β , IL-6 and TNF- α have been reported in mouse models ^{21,22} and a more generalised reduction in cytokine production in response to activation of TLR 2, 3, 4, 5, 6 and 9 was observed in macrophage from aged C57BL/6 mice ²³. Human studies of TLR function have also revealed age-associated declines in cytokine production with older subjects showing reductions in TLR1/2 induced IL-6 and TNF- α production in monocytes ²⁴. This functional defect in cytokine production was strongly correlated with a decrease in surface expression of TLR1 on the surface of monocytes from old compared with young individuals, and an age-associated decrease in TLR4 expression was also reported. Reduced TLR function and expression on monocytes and macrophage will compromise an individual's ability to both detect and eliminate harmful pathogens, thereby prolonging the duration and increasing the severity of infection.

Results from these studies would seem to contradict or question the observation of a heightened proinflammatory milieu in older individuals with higher circulating levels of cytokines such as IL-1 β , IL-6 and TNF- α and proinflammatory markers such as C-reactive protein commonly referred to as 'inflamm-ageing' ¹³. However, serum levels of cytokines might reflect production from a variety of tissue types. For example, although monocytes/macrophages are a major source of IL-6, it is also produced by endothelial cells, adipocytes, muscle cells, stromal cells and other cell types that are susceptible to the ageing process ²⁵.

Dendritic Cells (Antigen-Presenting Cells)

Antigen presentation is a critical event in the regulation of antigen-specific immune responses. Antigen-presenting cells (APCs) engulf, process and present pathogen-associated antigens on major histocompatibility complex (MHC) molecules recognised by T cells resulting in antigen-specific immune responses and long lived immunity ²⁶. APCs include macrophages, B cells and dendritic cells (DCs), however it is DCs that act as the main interface between the innate and adaptive immune system. DCs can be classified into two subsets: myeloid DCs (mDCs) and plasmacytoid DCs (pDCs) that are of a lymphoid lineage. Both mDCs and pDCs originate from a common DC precursor that originates from hematopoietic stem cells in the bone marrow ¹⁸. During pathogen invasion, DCs detect intruders via pattern recognition receptors (e.g. TLRs), capture antigens and quickly migrate to the draining lymph nodes where they will present antigen to T cells via MHC molecules and elicit a specific immune response.

One of the potential mechanisms by which ageing impacts the overall function of DCs is by alterations in number or distributions of this cell. However, there has been conflicting data reported to date. Significantly lower numbers of mDCs and pDCs have been observed in the frail elderly when compared to both healthy young and healthy elderly subjects ²⁷. Other studies have found no significant differences between peripheral mDCs and pDCs between young and elderly subjects ²⁸ while another observed a reduction in mDCs with age ²⁹. Studies have generally found impairments in inflammatory responses of DCs, principally to TLR activation. Reduced IFN responses of pDCs from aged individuals to both the influenza and West Nile virus have been reported ^{30,31}. This is of note as the influenza virus is known to induce higher mortality in older people ⁸ and older people exhibit more severe infection following contact with West Nile virus ³².

The Adaptive Immune System

B Cells

B cells are derived from pluripotent haematopoietic stem cells in the bone marrow. Their main function is the production of specific antibodies in response to a specific antigen and this function is critical for an effective response against extracellular bacterial infections and for vaccinations. High-affinity-specific antibodies are generated by somatic hypermutation of immunoglobulin (Ig) genes in the germinal centre of lymphoid tissue, after which professional antibody secreting plasma cells migrate to the bloodstream³³. Large amounts of antibodies are then secreted into circulation and provide defence against harmful pathogens through a number of processes. They can bind directly to antigens, effectively coating the surface of the invader, in order to prevent pathogens from entering or damaging healthy body cells. Antibodies can also stimulate other parts of the immune system (e.g. complement proteins) to destroy the pathogens. Furthermore, they can mark pathogens through a process known as opsonisation so that the pathogens can be identified and neutralised by other immune cells. Memory B cells are also generated, which provide immunological memory in case of subsequent infection with the same pathogen.



The total number of B cells has been reported to diminish with ageing³⁴. Additionally older people present with a decrease in the diversity of their B cell repertoire characterised by an increase in memory B cells and a decrease in naïve B cells^{35,36}. Consistent with the change in B cell numbers and subsets, antibody production is also affected by age. In response to vaccinations, older persons present a delayed response seen as a reduced clonal expansion of plasma cells. This finding correlates with decreases in Ig production and these Ig cells have lower affinities toward their antigens³⁷. Additionally, when a primary antibody response is required in the presence of novel antigens, the elderly display a delayed response with lower levels of high affinity antibodies³⁸. This would in part, explain the longer duration and increased severity of infections experienced by the elderly population.

T Cells

Most of the lymphocytes in the blood are T cells. While B cells mature in the bone marrow, T cell precursors migrate to and mature in the thymus. T cells are characterised by the presence of T-cell receptors (TCRs) and can be categorised into two main subsets by their surface expression of either CD4 or CD8⁶. CD4+ cells, which recognise antigens presented on class II major histocompatibility complex (MHC) are mainly regulatory cells, whereas CD8+ cells are cytotoxic and recognise antigen presented on class I MHC molecules. Both functions are of vital importance for both the adaptive and innate immune responses. CD4+ T cells are also referred to as T helper (Th) cells, as they assist other white blood cells in immunologic processes, including maturation of B cells into plasma cells and memory B cells, and activation of cytotoxic T cells and macrophages. Th cells can be further categorised into different subsets (e.g. Th1, Th2, Th17, Treg) based on production of signature cytokines⁷. CD8+ T cells (also known as cytotoxic T cells) are vital for immune defence against intracellular pathogens and for tumour surveillance. CD8+ T cells kill infected and malignant cells through the secretion of pro-inflammatory cytokines, the release of toxic granules called perforins and granzymes, and through the use of Fas/FasL interactions³⁹.

Similar to B cells, the total number of naïve T cells decreases with age. T cells develop in the thymus, which decreases with age due to age-related changes that affect both T-cell progenitors and the thymic microenvironment⁴⁰. This decline affects CD4+ and CD8+ cells differently with a better preserved population of naïve CD4+ cells and greater decline in CD8+ cells (Ferrando-Martinez et al. 2011). Alterations in the CD8+ T cell compartment are some of the best characterised age-related changes in the immune system. These cells tend to undergo oligoclonal expansion with age and, as such the CD8+ T cell repertoire becomes increasingly skewed towards previously encountered antigens^{42,43}. There is an association between the increase in the prevalence of CD8+ T cell oligoclonal expansion and cytomegalovirus (CMV) infection status, which establishes life-long latent infection⁴². This oligoclonal expansion can also limit the ability of the CD8+ T cell population to respond to newly encountered pathogens such as emerging strains of the influenza virus, which could explain the increased morbidity and mortality rates associated with influenza infection in the elderly. Activation of both naïve and memory T cells is a complex process that requires the intervention of co-stimulatory molecules such as CD28 after the binding of the TCR to MHC molecules⁴⁴. Binding of CD28 to its co-receptor results in a potent activation stimuli for T cells and although this activation is not impaired with ageing, the expression levels of CD28 in both CD4+ and CD8+ T cells decreases⁴⁵. This is consistent with a decreased naïve T cell pool and the accumulation of highly differentiated T cells.



Fiona McEvoy, Ph.D. is a postdoctoral researcher at Dublin City University. Her studies primarily focus on identifying novel compounds that can modulate the immune system to provide therapeutic benefits using in vitro and in vivo models.

Christine Loscher, Ph.D. is an Associate Professor of Immunology at Dublin City University. She is also the Lead Investigator of the Immunomodulation Research Group and has developed significant expertise in commercial research and industry engagement. She has secured over €4.5M in external funding (€400K directly from industry) for her research.

References:

1. United Nations, Department of Economic and Social Affairs, P. D. (2015). World Population Ageing 2015. *World Popul. Ageing 2015 (ST/ESA/SER.A/390)*.
2. Aw, D., Silva, A. B. & Palmer, D. B. Immunosenescence: Emerging challenges for an ageing population. *Immunology* **120**, 435-446 (2007).
3. Maijo, M., Clements, S. J., Ivory, K., Nicoletti, C. & Carding, S. R. Nutrition, diet and immunosenescence. *Mech. Ageing Dev.* **136-137**, 116-128 (2014).
4. Medzhitov, R. Recognition of microorganisms and activation of the immune response. *Nature* **449**, 819-26 (2007).
5. Kumar, H., Kawai, T. & Akira, S. Toll-like receptors and innate immunity. *Biochem. Biophys. Res. Commun.* **388**, 621-625 (2009).
6. Mosmann, T. R. & Sad, S. The expanding universe of T-cell subsets: Th1, Th2 and more. *Immunol. Today* **17**, 138-146 (1996).
7. Zhu, J., Yamane, H. & Paul, W. Differentiation of effector CD4 T cell populations. *Annu Rev Immunol.* **28**, 445-489 (2010).
8. Katz, J. M., Renshaw-hoelscher, M. & Tumpey, T. M. Immunity to Influenza. *Immunol. Res.* **29**, 113-124 (2004).
9. Weinberger, B., Herndler-Brandstetter, D., Schwanninger, A., Weiskopf, D. & Grubeck-Loebenstien, B. Biology of immune responses to vaccines in elderly persons. *Clin Infect Dis* **46**, 1078-1084 (2008).
10. Fulop, T. et al. Potential role of immunosenescence in cancer development. *Ann. N. Y. Acad. Sci.* **1197**, 158-165 (2010).
11. Prelog, M. Aging of the immune system: A risk factor for autoimmunity? *Autoimmun. Rev.* **5**, 136-139 (2006).
12. Krabbe, K. S., Pedersen, M. & Bruunsgaard, H. Inflammatory mediators in the elderly. *Exp. Gerontol.* **39**, 687-699 (2004).
13. Michaud, M. et al. Proinflammatory cytokines, aging and age-related diseases. *J. Am. Med. Disord. Assoc.* **14**, 877-882 (2013).
14. Shaw, A. C., Joshi, S., Greenwood, H., Panda, A. & Lord, J. M. Aging of the innate immune system. *Curr. Opin. Immunol.* **22**, 507-13 (2010).
15. Fortin, C. F., Larbi, A., Lesur, O., Douziech, N. & Fulop Jr., T. Impairment of SHP-1 down-regulation in the lipid rafts of human neutrophils under GM-CSF stimulation contributes to their age-related, altered functions. *J. Leukoc. Biol.* **79**, 1061-1072 (2006).
16. Wensch, C., Patruta, S., Daxböck, F., Krause, R. & Hörl, W. Effect of age on human neutrophil function. *J. Leukoc. Biol.* **67**, 40-45 (2000).
17. Fortin, C. F., Larbi, A., Dupuis, G., Lesur, O. & Fülöp, T. GM-CSF activates the Jak/STAT pathway to rescue polymorphonuclear neutrophils from spontaneous apoptosis in young but not elderly individuals. *Biogerontology* **8**, 173-187 (2007).
18. Geissman, F. et al. Development of monocytes, macrophages, and dendritic cells. *J. Microbiol. Immunol. Infect.* 656-662 (2010).
19. Gordon, S. & Taylor, P. R. Monocyte and macrophage heterogeneity. *Nat. Rev. Immunol.* **5**, 953-64 (2005).
20. Linehan, E. & Fitzgerald, D. C. Ageing and the immune system: focus on macrophages. *Eur. J. Microbiol. Immunol. (Bp).* **5**, 14-24 (2015).
21. Boehmer, E. D., Goral, J., Faunce, D. E. & Kovacs, E. J. Age-dependent decrease in Toll-like receptor 4-mediated proinflammatory cytokine production and mitogen- activated protein kinase expression. *J. Leukoc. Biol.* **75**, 342-349 (2004).
22. Chelvarajan, R. L., Collins, S. M., Van Willigen, J. M. & Bondada, S. The unresponsiveness of aged mice to polysaccharide antigens is a result of a defect in macrophage function. *J. Leukoc. Biol.* **77**, 503-12 (2005).
23. Renshaw, M. et al. Cutting edge: impaired Toll-like receptor expression and function in aging. *J. Immunol.* **169**, 4697-701 (2002).
24. Van Duin, H. et al. Function Age-Associated Defect in Human TLR-1/2 Age-Associated Defect in Human TLR-1/2 Function. *J Immunol Ref.* **178**, 970-975 (2007).
25. Maggio, D. et al. 25(OH)D Serum levels decline with age earlier in women than in men and less efficiently prevent compensatory hyperparathyroidism in older adults. *J. Gerontol. A. Biol. Sci. Med. Sci.* **60**, 1414-1419 (2005).
26. Guermonprez, P., Valladeau, J., Zitvogel, L., Théry, C. & Amigorena, S. ANTIGEN PRESENTATION AND T CELL STIMULATION BY DENDRITIC CELLS. *Annu. Rev. Immunol* **20**, 621-67 (2002).
27. Jing, Y. et al. Aging is associated with a numerical and functional decline in plasmacytoid dendritic cells, whereas myeloid dendritic cells are relatively unaltered in human peripheral blood. *Hum. Immunol.* **70**, 777-784 (2009).
28. Agrawal, A. et al. Altered Innate Immune Functioning of Dendritic Cells in Elderly Humans: A Role of Phosphoinositide 3-Kinase-Signaling Pathway. *J. Immunol.* **178**, 6912-6922 (2007).

29. Della Bella, S. et al. Peripheral blood dendritic cells and monocytes are differently regulated in the elderly. *Clin. Immunol.* **122**, 220-228 (2007).
30. Prakash, S., Agrawal, S., Cao, J., Gupta, S. & Agrawal, A. Impaired secretion of interferons by dendritic cells from aged subjects to influenza : role of histone modifications. *Age (Dordr).* **35**, 1785-97 (2013).
31. Qian, F. et al. Impaired interferon signaling in dendritic cells from older donors infected in vitro with West Nile virus. *J. Infect. Dis.* **203**, 1415-24 (2011).
32. Colpitts, T. M., Conway, M. J., Montgomery, R. R. & Fikrig, E. West Nile Virus: biology, transmission, and human infection. *Clin. Microbiol. Rev.* **25**, 635-48 (2012).
33. Hardy, R. R. & Hayakawa, K. B Cell development PATHways. *Annu. Rev. Immunol.* **19**, 595-621 (2001).
34. Ademokun, A., Wu, Y. C. & Dunn-Walters, D. The ageing B cell population: Composition and function. *Biogerontology* **11**, 125-137 (2010).
35. Gibson, K. L. et al. B-cell diversity decreases in old age and is correlated with poor health status. *Aging Cell* **8**, 18-25 (2009).
36. Chong, Y. et al. CD27+ (memory) B cell decrease and apoptosis-resistant CD27- (naive) B cell increase in aged humans: Implications for age-related peripheral B cell developmental disturbances. *Int. Immunol.* **17**, 383-390 (2005).
37. Sasaki, S. et al. Limited efficacy of inactivated influenza vaccine in elderly individuals is associated with decreased production of vaccine-specific antibodies. *J. Clin. Invest.* **121**, 3109-3119 (2011).
38. Roukens, A. H. et al. Elderly subjects have a delayed antibody response and prolonged viraemia following yellow fever vaccination: A prospective controlled cohort study. *PLoS One* **6**, 1-6 (2011).
39. Tschärke, D. C., Croft, N. P., Doherty, P. C. & La Gruta, N. L. Sizing up the key determinants of the CD8+ T cell response. *Nat. Rev. Immunol.* **15**, 705-716 (2015).
40. Aspinall, R. & Andrew, D. Thymic involution in aging. *J. Clin. Immunol.* **20**, 250-256 (2000).
41. Ferrando-Martínez, S. et al. Age-related deregulation of naive T cell homeostasis in elderly humans. *Age (Omaha).* **33**, 197-207 (2011).
42. Colonna-Romano, G. et al. Impact of CMV and EBV seropositivity on CD8 T lymphocytes in an old population from West-Sicily. *Exp. Gerontol.* **42**, 995-1002 (2007).
43. Akbar, A. N. & Fletcher, J. M. Memory T cell homeostasis and senescence during aging. *Curr. Opin. Immunol.* **17**, 480-485 (2005).
44. Mempel, T. R., Henrickson, S. E. & von Andrian, U. H. T-cell priming by dendritic cells in lymph nodes occurs in three distinct phases. *Nature* **427**, 154-159 (2004).
45. Weng, N., Akbar, A. N. & Goronzy, J. CD28- T cells: their role in the age-associated decline of immune function. *Trends Immunol.* **30**, 306-312 (2009).
46. Franceschi, C. & Campisi, J. Chronic inflammation (Inflammaging) and its potential contribution to age-associated diseases. *Journals Gerontol. - Ser. A Biol. Sci. Med. Sci.* **69**, S4-S9 (2014).
47. Trichopoulou, A. Traditional Mediterranean diet and longevity in the elderly: a review. *Public Health Nutr.* **7**, 943-947 (2004).
48. Brambilla, D. et al. The role of antioxidant supplement in immune system, neoplastic, and neurodegenerative disorders: a point of view for an assessment of the risk/benefit profile. *Nutr. J.* **7**, 29 (2008).
49. Meydani, S. N. et al. Vitamin E supplementation enhances cell-mediated immunity in healthy elderly subjects. *Am. J. Clin. Nutr.* **52**, 557-563 (1990).
50. Meydani, S. N. et al. Vitamin E supplementation and in vivo immune response in healthy elderly subjects. A randomized controlled trial. *JAMA* **277**, 1380-6 (1997).
51. De la Fuente, M., Hernanz, A., Guayerbas, N., Victor, V. M. & Arnalich, F. Vitamin E ingestion improves several immune functions in elderly men and women. *Free Radic. Res.* **42**, 272-80 (2008).
52. Wu, D. & Meydani, S. N. Age-associated changes in immune and inflammatory responses: impact of vitamin E intervention. *J. Leukoc. Biol.* **84**, 900-14 (2008).
53. Hayek, M. G. et al. Vitamin E supplementation decreases lung virus titers in mice infected with influenza. *J. Infect. Dis.* **176**, 273-6 (1997).

54. Han, S. N. et al. Vitamin E supplementation increases T helper 1 cytokine production in old mice infected with influenza virus. *Immunology* **100**, 487–493 (2000).
55. Chavance, M., Herbeth, B., Fournier, C., Janot, C. & Vernhes, G. Vitamin status, immunity and infections in an elderly population. *Eur. J. Clin. Nutr.* **43**, 827–35 (1989).
56. Meydani, S. N., Han, S. N. & Wu, D. Vitamin E and immune response in the aged: Molecular mechanisms and clinical implications. *Immunol. Rev.* **205**, 269–284 (2005).
57. Battault, S. et al. Vitamin D metabolism, functions and needs: From science to health claims. *Eur. J. Nutr.* **52**, 429–441 (2013).
58. Borel, P., Caillaud, D. & Cano, N. J. Vitamin D bioavailability: state of the art. *Crit. Rev. Food Sci. Nutr.* **55**, 1193–205 (2015).
59. Hilger, J. et al. A systematic review of vitamin D status in populations worldwide. *Br. J. Nutr.* **111**, 23–45 (2014).
60. Prietl, B., Treiber, G., Pieber, T. R. & Amrein, K. Vitamin D and immune function. *Nutrients* **5**, 2502–2521 (2013).
61. Sadeghi, K. et al. Vitamin D3 down-regulates monocyte TLR expression and triggers hyporesponsiveness to pathogen-associated molecular patterns. *Eur. J. Immunol.* **36**, 361–370 (2006).
62. Calton, E. K., Keane, K. N., Newsholme, P. & Soares, M. J. The impact of Vitamin D levels on inflammatory status: A systematic review of immune cell studies. *PLoS One* **10**, 1–12 (2015).
63. Perlstein, T. S., Pande, R., Berliner, N. & Vanasse, G. J. Prevalence of 25-hydroxyvitamin D deficiency in subgroups of elderly persons with anemia: Association with anemia of inflammation. *Blood* **117**, 2800–2806 (2011).
64. Laird, E. et al. Vitamin D deficiency is associated with inflammation in older Irish adults. *J. Clin. Endocrinol. Metab.* **99**, 1807–1815 (2014).
65. Tang, B. M., Eslick, G. D., Nowson, C., Smith, C. & Bensoussan, A. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet* **370**, 657–666 (2007).
66. Schrezenmeir, J. & de Vrese, M. Probiotics, prebiotics, and synbiotics—approaching a definition. *Am. J. Clin. Nutr.* **73**, 361S–364S (2001).
67. Biagi, E., Candela, M., Fairweather-Tait, S., Franceschi, C. & Brigidi, P. Aging of the human metaorganism: the microbial counterpart. *Age (Dordr.)* **34**, 247–67 (2012).
68. Man, A. L., Gicheva, N. & Nicoletti, C. The impact of ageing on the intestinal epithelial barrier and immune system. *Cell. Immunol.* **289**, 112–118 (2014).
69. Fujihashi, K. & Kiyono, H. Mucosal immunosenescence: new developments and vaccines to control infectious diseases. *Trends Immunol.* **30**, 334–343 (2009).
70. Yan, F. & Polk, D. B. Probiotics and immune health. *Curr. Opin. Gastroenterol.* **27**, 496–501 (2011).
71. Maneerat, S. et al. Consumption of Bifidobacterium lactis Bi-07 by healthy elderly adults enhances phagocytic activity of monocytes and granulocytes. *J. Nutr. Sci.* **2**, (2013).
72. Boge, T. et al. A probiotic fermented dairy drink improves antibody response to influenza vaccination in the elderly in two randomised controlled trials. *Vaccine* **27**, 5677–5684 (2009).
73. Guillemard, E., Tondou, F., Lacoïn, F. & Schrezenmeir, J. Consumption of a fermented dairy product containing the probiotic Lactobacillus casei DN-114 001 reduces the duration of respiratory infections in the elderly in a randomised controlled trial. *Br. J. Nutr.* **103**, 58 (2010).
74. Makino, S. et al. Reducing the risk of infection in the elderly by dietary intake of yoghurt fermented with Lactobacillus delbrueckii ssp. bulgaricus OLL1073R-1. *Br. J. Nutr.* **104**, 998–1006 (2010).
75. Fritsche, K. Fatty acids as modulators of the immune response. *Annu. Rev. Nutr.* **26**, 45–73 (2006).
76. Calder, P. C. The 2008 ESPEN Sir David Cuthbertson lecture: Fatty acids and inflammation - From the membrane to the nucleus and from the laboratory bench to the clinic. *Clin. Nutr.* **29**, 5–12 (2010).
77. Pae, M., Meydani, S. N. & Wu, D. The role of nutrition in enhancing immunity in aging. *Aging Dis.* **3**, 91–129 (2012).
78. Rees, D. et al. Dose-related effects of eicosapentaenoic acid on innate immune function in healthy humans: a comparison of young and older men. *Am. J. Clin. Nutr.* **83**, 331–342 (2006).
79. Draper, E. et al. Omega-3 fatty acids attenuate dendritic cell function via NF-κB independent of PPARγ. *J. Nutr. Biochem.* **22**, 784–790 (2011).

80. Weldon, S. M., Mullen, A. C., Loscher, C. E., Hurley, L. A. & Roche, H. M. Docosahexaenoic acid induces an anti-inflammatory profile in lipopolysaccharide-stimulated human THP-1 macrophages more effectively than eicosapentaenoic acid. *J. Nutr. Biochem.* **18**, 250–258 (2007).
81. Meydani, S. N. et al. Oral (n-3) fatty acid supplementation suppresses cytokine production and lymphocyte proliferation: comparison between young and older women. *J. Nutr.* **121**, 547–55 (1991).
82. Bechoua, S. et al. Influence of very low dietary intake of marine oil on some functional aspects of immune cells in healthy elderly people. *Br. J. Nutr.* **89**, 523–31 (2003).
83. Bouwens, M. et al. Fish-oil supplementation induces antiinflammatory gene expression profiles in human blood mononuclear cells. *Am J Clin Nutr* **90**, 415–424 (2009).
84. Alperovich, M., Neuman, M. I., Willett, W. C. & Curhan, G. C. Fatty acid intake and the risk of community-acquired pneumonia in U.S. women. *Nutrition* **23**, 196–202 (2007).
85. Merchant, A. T., Curhan, G. C., Rimm, E. B., Willett, W. C. & Fawzi, W. W. Intake of n6 and n3 fatty acids and fish and risk of community- acquired pneumonia in US men 1 – 3. *Am. J. Clin. Nutr.* **82**, 668–674 (2005).
86. Boirie, Y., Morio, B., Caumon, E. & Cano, N. J. Nutrition and protein energy homeostasis in elderly. *Mech. Ageing Dev.* **136-137**, 76–84 (2014).
87. Gryson, C. et al. 'Fast proteins' with a unique essential amino acid content as an optimal nutrition in the elderly: Growing evidence. *Clin. Nutr.* **33**, 642–648 (2014).
88. Allison, S. P. Malnutrition, disease, and outcome. *Nutrition* **16**, 590–593 (2000).
89. Milne, a. C., Potter, J., Vivanti, a & Avenell, a. Protein and energy supplementation in older people at risk from malnutrition (2009). *Australas. J. Ageing* **29**, 144 (2010).
90. Marshall, K. Therapeutic applications of whey protein. *Altern. Med. Rev.* **9**, 136–156 (2004).
91. Brandenburg, K., Heinbockel, L., Correa, W. & Lohner, K. Peptides with dual mode of action: Killing bacteria and preventing endotoxin-induced sepsis. *Biochim. Biophys. Acta - Biomembr.* **1858**, 971–979 (2016).
92. Chatterton, D. E. W., Nguyen, D. N., Bering, S. B. & Sangild, P. T. Anti-inflammatory mechanisms of bioactive milk proteins in the intestine of newborns. *Int. J. Biochem. Cell Biol.* **45**, 1730–1747 (2013).
93. Rusu, D., Drouin, R., Pouliot, Y., Gauthier, S. & Poubelle, P. E. A bovine whey protein extract can enhance innate immunity by priming normal human blood neutrophils. *J. Nutr.* **139**, 386–393 (2009).
94. Rusu, D., Drouin, R., Pouliot, Y., Gauthier, S. & Poubelle, P. E. A bovine whey protein extract stimulates human neutrophils to generate bioactive IL-1Ra through a NF-kappaB- and MAPK-dependent mechanism. *J. Nutr.* **140**, 382–91 (2010).
95. Chevalley, T., Hoffmeyer, P., Bonjour, J. P. & Rizzoli, R. Early serum IGF-I response to oral protein supplements in elderly women with a recent hip fracture. *Clin. Nutr.* **29**, 78–83 (2010).
96. Pal, S. & Ellis, V. The chronic effects of whey proteins on blood pressure, vascular function, and inflammatory markers in overweight individuals. *Obesity (Silver Spring)*. **18**, 1354–1359 (2010).
97. Lee, Y. M., Skurk, T., Hennig, M. & Hauner, H. Effect of a milk drink supplemented with whey peptides on blood pressure in patients with mild hypertension. *Eur. J. Nutr.* **46**, 21–27 (2007).
98. Gouni-Berthold, I. et al. The whey fermentation product malleable protein matrix decreases TAG concentrations in patients with the metabolic syndrome: a randomised placebo-controlled trial. *Br. J. Nutr.* **107**, 1694–706 (2012).
99. Petyaev, I. M., Dovgalevsky, P. Y., Klochkov, V. A., Chalyk, N. E. & Kyle, N. Whey protein lysosome formulation improves vascular functions and plasma lipids with reduction of markers of inflammation and oxidative stress in prehypertension. *ScientificWorldJournal*. **2012**, 269476 (2012).
100. Bharadwaj, S., Naidu, T. A. G., Betageri, G. V., Prasadarao, N. V. & Naidu, A. S. Inflammatory responses improve with milk ribonuclease-enriched lactoferrin supplementation in postmenopausal women. *Inflamm. Res.* **59**, 971–978 (2010).
101. Zhou, L. M. et al. Effect of whey supplementation on circulating C-reactive protein: A meta-analysis of randomized controlled trials. *Nutrients* **7**, 1131–1143 (2015).